

Sigma-Aldrich®

Lab & Production Materials

Merck



ELISA Analysis Kits

- Over 1,800 ELISAs now available
- Detection of cytokine markers for all species within our portfolio
- Validation criteria including: inter- and intra-assay CV, sensitivity, recovery linearity, biological range, cross-reactivity, species-specificity, sample and shipping stability, and method comparison
- Variability avoidance with accurate reagent dispensing and thorough plate washing
- All kits backed by our technical resources

To search, please visit:

https://www.scientificlabs.co.uk/ content/DEFAULT/1/lifescience/1604

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.



1/4

(@) slsinfo@scientific-labs.com scientificlabs.co.uk

00408



Our ELISA portfolio

Covering a diversity of research areas and species, our pre-validated, easy to use ELISAs deliver reliable and reproducible results for quantifying biomarkers in complex sample types at the pg/mL level. Our substantial portfolio currently includes over 1,800 sensitive and specific ELISAs - the easiest way to search them is at scientificlabs.co.uk

Research areas covered



Cancer



Toxicity



Neuroscience



Bone Metabolism



Cardiovascular **Disease**



Immunology

- Acute Phase Proteins
- Adipokines
- · Angiogenesis Biomarkers
- Apolipoproteins
- Apoptosis Pathway Proteins
- BMPs
- Bone Metabolism Biomarkers
- Cancer Biomarkers
- Cardiovascular Disease Biomarkers
- CD Markers
- Cell Signaling Pathway Proteins
- Cell Structure Proteins
- Chemokines
- Complement Factors
- Cytokines

- · Growth Factors
- Gut Hormones
- IGFBPs
- Immune Checkpoint Proteins
- · Immunoglobulins
- Kidney Injury Biomarkers
- Liver Proteins
- · Metabolic Hormones
- MMPs
- Neurodegenerative Disease Biomarkers
- Neurological Disorders Biomarkers
- · Sepsis Biomarkers
- Soluble Receptors
- Thyroid Hormones
- · Vascular Injury Biomarkers

Species within our portfolio

| | Cytokines | Metabolic | Cardiovascular | Angiogenesis | Signaling |
|---------------------------|-----------|-----------|----------------|--------------|-----------|
| | ✓ | V | V | V | V |
| | ✓ | ✓ | ✓ | ~ | V |
| - | ✓ | V | | V | V |
| | ✓ | | | | |
| Ai | ✓ | V | | | |
| | ✓ | | | | |
| | V | | | | |
| | ✓ | | | | |
| a | ✓ | | | | |
| (All parties and a second | ✓ | ✓ | | | |







Metabolic/Endocrine ELISAs

Our menu of metabolic and endocrine ELISAs allows end-users a consistent and reliable method of investigating metabolic disease states or defining outcomes within toxicological studies during drug development.

| Description | Species | Standard Curve Range | Sensitivity | Sample Volume | Cat. No. | Bulk Packaging Cat. No.* |
|--|-------------------|-------------------------|-------------|------------------|-------------------|-----------------------------|
| Adiponectin | Human | 1.56-200 ng/mL | 0.2 ng/mL | 10 μL | EZHADP-61K | EZHADP-61BK |
| Adiponectin (High Molecular Weight) | Human | 1.5-200 ng/mL | 0.5 ng/mL | 10 µL | EZHMWAN-65K | |
| Adiponectin | Mouse | 1-50 ng/mL | 0.2 ng/mL | 10 µL | EZMADP-60K | EZMADP-60BK |
| Adiponectin | Rat | 3.125-200 ng/mL | 0.4 ng/mL | 10 µL | EZRADP-62K | |
| Amylin (active) | Human | 1-100 pM | 0.7 pM | 50 μL | EZHA-52K | |
| C-Peptide | Human | 0.2-20 ng/mL | 0.05 ng/mL | 10 µL | EZHCP-20K | EZHCP-20BK |
| C-Peptide | Canine | 0.2-10 ng/mL | 0.24 ng/mL | 25 µL | EZCCP-47K | |
| C-Peptide 2 | Mouse, Rat | 25-1600 pM | 15.0 pM | 20 µL | EZRMCP2-21K | |
| C-Reactive Protein (CRP) | Human | 0.12-10 ng/mL | 0.2 ng/mL | 2 μL | CYT298 | |
| C-Reactive Protein (CRP) | Rat | 4.9-10 ng/mL | 2.5 ng/mL | 2 μL | CYT294 | |
| FGF-21 | Human | 31.25-2000 pg/mL | 10.0 pg/mL | 50 μL | EZHFGF21-19K | |
| FGF-21 | Mouse, Rat | 49.4-12,000 pg/mL | 10.0 pg/mL | 10 µL | EZRMFGF21- 26K | |
| FGF-23 | Human | 9.9-2400 pg/mL | 3.5 pg/mL | 50 μL | EZHFGF23-32K | EZHFGF23- 32BK |
| Ghrelin (active) | Human | 25-2000 pg/mL | 15.0 pg/mL | 20 µL | EZGRA-88K | EZGRA-88BK |
| Ghrelin (active) | Mouse, Rat | 25-2000 pg/mL | 8.0 pg/mL | 20 µL | EZRGRA-90K | |
| Ghrelin (total) | Human | 100-5000 pg/mL | 50.0 pg/mL | 20 µL | EZGRT-89K | EZGRT-89BK |
| Ghrelin (total) | Mouse, Rat | 0.1-10 ng/mL | 0.04 ng/mL | 20 µL | EZRGRT-91K | |
| GIP (total) | Human | 8.2-2000 pg/mL | 4.2 pg/mL | 20 µL | EZHGIP-54K | EZHGIP-54BK |
| GIP (total) | Mouse, Rat | 8.2-2000 pg/mL | 4.2 pg/mL | 10 µL | EZRMGIP-55K | EZRMGIP-55BK |
| GLP-1 (active) | Multi-Species | 2-100 pM | 1.0 pM | 100 μL | EGLP-35K | EGLP-35BK |
| GLP-1 High Sensitivity (active) Δ•• | Multi-Species | See data sheet | 0.14 pM | 50 μL | EZGLPHS-35K | EZGLPHS-35BK |
| GLP-1 (total) | Multi-Species | 4.1-1000 pM | 1.0 pM | 20-50 μL | EZGLP1T-36K | EZGLP1T-36BK |
| GLP-2 | Multi-Species | 1-64 ng/mL | 0.3 ng/mL | 50 μL | EZGLP2-37K | |
| Glucagon Δ | Human, Mouse, Rat | 0.02-2 ng/mL | 0.003 ng/mL | 150-300 μL | EZGLU-30K | EZGLU-30BK |
| Growth Hormone (GH) | Mouse, Rat | 0.7-50 ng/mL | 0.07 ng/mL | 10 μL | EZRMGH-45K | EZRMGH-45BK |
| Insulin | Human | 2-200 µU/mL | 1.0 μU/mL | 20 µL | EZHI-14K | EZHI-14BK |
| Insulin | Mouse, Rat | 0.2-10 ng/mL | 0.1 ng/mL | 10 µL | EZRMI-13K | EZRMI-13BK |
| Insulin (Animal serum free) | Human | 2-200 µU/mL | 0.85 μU/mL | 20 µL | EZHIASF-14K | |
| Leptin | Canine | 0.78-50 ng/mL | 0.21 ng/mL | 20 µL | EZCL-31K | |
| Leptin | Mouse | 0.2-30 ng/mL | 0.05 ng/mL | 10 µL | EZML-82K | EZML-82BK |
| Leptin | Rat | 0.2-30 ng/mL | 0.08 ng/mL | 10 µL | EZRL-83K | EZRL-83BK |
| Leptin "Dual Range" | Human | 0.5-100 ng/mL | 0.2 ng/mL | 25 µL | EZHL-80SK | EZHL-80BK |
| Procollagen Type IIA N-Propeptide (PIIANP) | Human | lot dependent | 30.0 ng/mL | 5 μL | EZPIIANP-53K | |
| Proinsulin (total) | Human | 2-200 pM | 0.5 pM | 20 µL | EZHPI-15K | EZHPI-15BK |
| PYY (total) | Human | 10-2000 pg/mL | 6.5 pg/mL | 20 µL | EZHPYYT66K | |
| Resistin | Human | 0.16-5 ng/mL | 0.02 ng/mL | 20 µL | EZHR-95K | EZHR-95BK |
| SAA-3 | Mouse | 0.078-5 μg/mL | 0.078 μg/mL | 10 µL | EZMSAA3-12K | |

Bulk packaging available on select kits — more environmentally friendly and saves space (10 kit equivalent)

https://www.scientificlabs.co.uk/content/DEFAULT/1/lifescience/1604







Preferred assay for measuring GLP-1 (active)

Chemiluminescent assay



Neuroscience ELISAs

Get a complete picture of the complexities associated with normal and disease states of the nervous system with reliable quantification of biomarkers using our neuropeptide and neurodegenerative ELISAs.

| Description | Species | Standard Curve Range | Sensitivity | Sample Volume | Cat. No. |
|--|-------------------|--|--------------|------------------|--------------|
| Amyloid beta 1-40 | Human | 16-500 pg/mL | 4 pg/mL | 50 μL | EZHS40 |
| Amyloid beta 1-42 | Human | 16-500 pg/mL | 5 pg/mL | 50 μL | EZHS42 |
| Amyloid beta, Set | Human | Contains 1 each of EZHS40 and EZHS42 | | | EZHS-SET |
| Amyloid beta (Brain) 1-40 | Human | 16-500 pg/mL | 4 pg/mL | 50 μL | EZBRAIN40 |
| Amyloid beta (Brain) 1-42 | Human | 16-500 pg/mL | 5 pg/mL | 50 μL | EZBRAIN42 |
| Amyloid beta (Brain), Set | Human | Contains 1 each of EZBRAIN40 and EZBRAIN42 | | | EZBRAIN-SET |
| BDNF (Brain-Derived Neurotrophic Factor) | Human, Rat | 7.8-500 pg/mL | 7.8 pg/mL | 50 µL | CYT306 |
| GFAP (Glial Fibrillary Acidic Protein) | Human, Mouse, Rat | 1.5-100 ng/mL | 1.5 ng/mL | 100 µL | NS830 |
| NPY (Neuropeptide Y) | Human | 5-1000 pg/mL | 2 pg/mL | 50 μL | EZHNPY-25K |
| NPY (Neuropeptide Y) | Mouse, Rat | 0.01-2 ng/mL | 0.004 ng/mL | 20 μL | EZRMNPY-27K |
| PEDF (Pigment Epithelium- Derived Factor) | Human | 0.9-62.5 ng/mL | 0.9 ng/mL | 50 μL | CYT420 |
| Phosphorylated Neurofilament, (pNF-H) Sandwich | Multi-Species | 0.0293-15 ng/mL | 0.0585 ng/mL | 1-10 µL | NS170 |
| S100B | Human | 2.7-2000 pg/mL | 1.3 pg/mL | 50 μL | EZHS100B-33K |

Radioimmunoassays (RIAs)

RIAs have long been considered a valuable, economical and accurate way to measure protein concentration. As many researchers move away from using radioactive material, please consider using our other high quality immunoassays for your research needs.

| Description | Species | Standard Curve Range | Sensitivity | Sample Volume | Cat. No. |
|---------------------|---------------|-------------------------|-------------|------------------|------------|
| Adiponectin | Human | lot dependent | 1 ng/mL | 5 μL | HADP-61HK |
| C-Peptide | Canine | 0.156-20 ng/mL | 0.15 ng/mL | 50 μL | CCP-24HK |
| C-Peptide | Human | 0.1-5 ng/mL | 0.1 ng/mL | 50 μL | HCP-20K |
| Ghrelin (active) | Human | lot dependent | 7.8 pg/mL | 50 µL | GHRA-88HK |
| Ghrelin (total) | Human | lot dependent | 93 pg/mL | 50 μL | GHRT-89HK |
| GLP-1 (active) | Multi-Species | 10-500 pM | 3 pM | 300 µL | GLP1A-35HK |
| GLP-1 (total) | Multi-Species | 10-1000 pM | 3 pM | 300 µL | GLP1T-36HK |
| Glucagon | Multi-Species | 20-400 pg/mL | 20 pg/mL | 50 μL | GL-32K |
| Insulin | Porcine | 2-200 µU/mL | 2 μU/mL | 50 μL | PI-12K |
| Insulin | Rat | 0.1-10 ng/mL | 0.1 ng/mL | 50 μL | RI-13K |
| Insulin LisPro | Multi-Species | 2.5-250 μU/mL | 2.5 μU/mL | 100 µL | LPI-16K |
| Insulin (sensitive) | Rat | 0.02-1 ng/mL | 0.02 ng/mL | 50 μL | SRI-13K |
| Insulin Specific | Human | 2-200 μU/mL | 2 μU/mL | 50 μL | HI-14K |
| Leptin | Human | 0.5-100 ng/mL | 0.5 ng/mL | 50-100 μL | HL-81K |
| Leptin | Multi-Species | 1-50 ng/mL | 1 ng/mL | 50 μL | XL-85K |
| Proinsulin | Human | 2-100 pM | 2 pM | 100-200 μL | HPI-15K |
| PYY (3-36) | Human | lot dependent | 20 pg/mL | <100 µL | PYY-67HK |
| PYY (total) | Human | lot dependent | 10 pg/mL | <100 µL | РҮҮТ-66НК |

https://www.scientificlabs.co.uk/content/DEFAULT/1/lifescience/1604

© 2020 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved. Merck, Millipore, the vibrant M, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources. 32389 07/2020







